

# Interactive Example Candidate Responses

## Paper 4 (May/June 2016), Question 1

### Cambridge International AS & A Level Physics 9702

In order to help us develop the highest quality resources, we are undertaking a continuous programme of review; not only to measure the success of our resources but also to highlight areas for improvement and to identify new development needs.

We invite you to complete our survey by visiting the website below. Your comments on the quality and relevance of our resources are very important to us.

[www.surveymonkey.co.uk/r/GL6ZNJB](http://www.surveymonkey.co.uk/r/GL6ZNJB)

Would you like to become a Cambridge International consultant and help us develop support materials?

Please follow the link below to register your interest.

[www.cambridgeinternational.org/cambridge-for/teachers/teacherconsultants/](http://www.cambridgeinternational.org/cambridge-for/teachers/teacherconsultants/)

Copyright © UCLES 2018

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

UCLES retains the copyright on all its publications. Registered Centres are permitted to copy material from this booklet for their own internal use. However, we cannot give permission to Centres to photocopy any material that is acknowledged to a third party, even for internal use within a Centre.

- 1 A binary star consists of two stars A and B that orbit one another, as illustrated in Fig. 1.1.




Fig. 1.1

The stars are in circular orbits with the centres of both orbits at point P, a distance  $d$  from the centre of star A.

- (a) (i) Explain why the centripetal force acting on both stars has the same magnitude.

Because the centripetal force acting on both stars are provided by the gravitational force  $F_g = \frac{M_A M_B}{G r^2} = m \omega^2 r$   
 $= m \left( \frac{2\pi}{T} \right)^2 r$  the angular velocity  $\omega$  and period  $T$  of both stars are the same. So the centripetal force (and gravitational forces) for both stars are the same.

- (ii) The period of the orbit of the stars about point P is 4.0 years. Calculate the angular speed  $\omega$  of the stars.

$$\omega = \frac{2\pi}{T} = \frac{2\pi}{4 \times 365 \times 24 \times 3600}$$

$$= 4.98 \times 10^{-8} \text{ rad s}^{-1}$$

$$\omega = 4.98 \times 10^{-8} \text{ rad s}^{-1} \quad [2]$$

Your  
Mark

1(a)(i)

1(a)(ii)

1(b)(i)

1(b)(ii)

Q1	Mark scheme	
(a)(i)	gravitational force provides/is the centripetal force same gravitational force (by Newton III)	B1 B1 [2]
(a)(ii)	$\omega = 2\pi / T$ $= 2\pi / (4.0 \times 365 \times 24 \times 3600)$ $= 5.0 \text{ (4.98)} \times 10^{-8} \text{ rad s}^{-1}$	C1 A1 [2]
(b)(i)	(centripetal force =) $M_A d \omega^2 = M_B (2.8 \times 10^8 - d) \omega^2$ or $M_A d_A = M_B d_B$ $M_A / M_B = 3.0 = (2.8 \times 10^8 - d) / d$ $d = 7.0 \times 10^7 \text{ km}$	C1 C1 A1 [3]
(b)(ii)	$G M_A M_B / (2.8 \times 10^{11})^2 = M_A d \omega^2$ $M_B = (2.8 \times 10^{11})^2 \times d \omega^2 / G$ $= (2.8 \times 10^{11})^2 \times (7.0 \times 10^{10}) \times (4.98 \times 10^{-8})^2 /$ $(6.67 \times 10^{-11})$ $= 2.0 \times 10^{29} \text{ kg}$	B1 C1 A1 [3]
		[Total: 10]

- (b) The separation of the centres of the stars is  $2.8 \times 10^8 \text{ km}$ .  
The mass of star A is  $M_A$ . The mass of star B is  $M_B$ .  
The ratio  $\frac{M_A}{M_B}$  is 3.0.

- (i) Determine the distance  $d$ .

$\therefore \omega, T, F_c$  are the same for A and B  
 $\therefore M_A \omega^2 d = M_B \omega^2 (2.8 \times 10^8 - d)$

$$\therefore \frac{M_A}{M_B} = \frac{2.8 \times 10^8 - d}{d} = 3$$

$$\therefore d = 7.0 \times 10^7 \text{ km}$$

$$d = 7.0 \times 10^7 \text{ km [3]}$$

- (ii) Use your answers in (a)(ii) and (b)(i) to determine the mass  $M_B$  of star B.  
Explain your working.

$$\therefore \frac{M_A}{M_B} = 3 \quad \therefore M_A = 3M_B$$

$$\begin{aligned} (7 \times 10^7) M_A &= 3M_B \times 7 \times 10^7 = M_B \times (2.8 \times 10^8 - 7 \times 10^7) \\ L = d_A + d_B &\rightarrow d_B = L - d_A \quad d_A = L - d_B \\ M_A \cdot d_A &= M_B \cdot d_B \end{aligned}$$

$$\begin{aligned} M_A d_A &= M_B d_B \\ M_A (L - d_B) &= M_B d_B \\ M_B &= \frac{M_A (L - d_B)}{d_B} \end{aligned}$$

$$\therefore \frac{M_A}{M_B} = 3 \quad \therefore M_A = 3M_B$$

$$\therefore G \frac{3M_B \cdot M_B}{L^2} = M_B \omega^2 L$$

$$6.67 \times 10^{-11} \times \frac{3M_B}{(2.8 \times 10^8 \times 10^3)^2} = (4.98 \times 10^{-8})^2 \times (2.8 \times 10^8 \times 10^3 - 7 \times 10^7)$$

$$M_B = 2.72 \times 10^{29} \text{ kg}$$

Your  
Mark

1(a)(i)


1(a)(ii)

1(b)(i)

1(b)(ii)

Q1	Mark scheme	
(a)(i)	gravitational force provides/is the centripetal force same gravitational force (by Newton III)	B1 B1 [2]
(a)(ii)	$\omega = 2\pi / T$ $= 2\pi / (4.0 \times 365 \times 24 \times 3600)$ $= 5.0 (4.98) \times 10^{-8} \text{ rad s}^{-1}$	C1 A1 [2]
(b)(i)	(centripetal force =) $M_A d \omega^2 = M_B (2.8 \times 10^8 - d) \omega^2$ or $M_A d_A = M_B d_B$ $M_A / M_B = 3.0 = (2.8 \times 10^8 - d) / d$ $d = 7.0 \times 10^7 \text{ km}$	C1 C1 A1 [3]
(b)(ii)	$GM_A M_B / (2.8 \times 10^{11})^2 = M_A d \omega^2$ $M_B = (2.8 \times 10^{11})^2 \times d \omega^2 / G$ $= (2.8 \times 10^{11})^2 \times (7.0 \times 10^{10}) \times (4.98 \times 10^{-8})^2 / (6.67 \times 10^{-11})$ $= 2.0 \times 10^{29} \text{ kg}$	B1 C1 A1 [3]
		[Total: 10]

- 1 A binary star consists of two stars A and B that orbit one another, as illustrated in Fig. 1.1.



The stars are in circular orbits with the centres of both orbits at point P, a distance  $d$  from the centre of star A.

- (a) (i) Explain why the centripetal force acting on both stars has the same magnitude.

Both stars have same angular speed and acceleration. They act as point masses so mass has negligible effect on force. [2]

- (ii) The period of the orbit of the stars about point P is 4.0 years.

Calculate the angular speed  $\omega$  of the stars.

$$\omega = \frac{2\pi}{T} = \frac{2\pi f}{1} = \frac{2\pi}{4 \times 365 \times 24 \times 3600}$$

$$\omega = 4.98 \times 10^{-8} \text{ rad s}^{-1} \quad [2]$$

Your  
Mark

1(a)(i)

1(a)(ii)

1(b)(i)

1(b)(ii)

Q1	Mark scheme	
(a)(i)	gravitational force provides/is the centripetal force same gravitational force (by Newton III)	B1 B1 [2]
(a)(ii)	$\omega = 2\pi / T$ $= 2\pi / (4.0 \times 365 \times 24 \times 3600)$ $= 5.0 \text{ (4.98)} \times 10^{-8} \text{ rad s}^{-1}$	C1 A1 [2]
(b)(i)	(centripetal force =) $M_A d \omega^2 = M_B (2.8 \times 10^8 - d) \omega^2$ or $M_A d_A = M_B d_B$ $M_A / M_B = 3.0 = (2.8 \times 10^8 - d) / d$ $d = 7.0 \times 10^7 \text{ km}$	C1 C1 A1 [3]
(b)(ii)	$GM_A M_B / (2.8 \times 10^{11})^2 = M_A d \omega^2$ $M_B = (2.8 \times 10^{11})^2 \times d \omega^2 / G$ $= (2.8 \times 10^{11})^2 \times (7.0 \times 10^{10}) \times (4.98 \times 10^{-8})^2 / (6.67 \times 10^{-11})$ $= 2.0 \times 10^{29} \text{ kg}$	B1 C1 A1 [3]
		[Total: 10]

Your  
Mark

1(a)(i)

1(a)(ii)

1(b)(i)

1(b)(ii)

- (b) The separation of the centres of the stars is  $2.8 \times 10^8$  km.  
The mass of star A is  $M_A$ . The mass of star B is  $M_B$ .

The ratio  $\frac{M_A}{M_B}$  is 3.0.

$F_c$  force.

- (i) Determine the distance  $d$ .

$$M_A \times d \times \omega^2 = M_B \times (2.8 \times 10^8 - d) \times \omega^2$$

$$\frac{M_A}{M_B} \times d = (2.8 \times 10^8 - d)$$

$$3 \times d = 2.8 \times 10^8 - d$$

$$3d + d = 2.8 \times 10^8$$

$$4d = 2.8 \times 10^8 \quad d = 7.0 \times 10^7 \text{ km [3]}$$

- (ii) Use your answers in (a)(ii) and (b)(i) to determine the mass  $M_B$  of star B.  
Explain your working.

$$F_c = m r \omega^2$$

$$r = 2.8 \times 10^8 \text{ km}$$

$$M_A \omega^2 = \frac{G M_A}{r^2}$$

$$(7.0 \times 10^7)^2 \times (4.98 \times 10^{-8})^2 = \frac{(6.67 \times 10^{-11})(M_A)}{(7.0 \times 10^7)^2}$$

$$\frac{M_A}{M_B} = 3 \quad \begin{matrix} 2602.7 \\ 2600 \end{matrix} \quad 2.6 \times 10^3 \text{ kg}$$

$$M_B = 8.7 \times 10^2 \text{ kg [3]}$$

$$\frac{(2.6 \times 10^3)}{3} = M_B \quad 876$$

[Total: 10]

Q1	Mark scheme	
(a)(i)	gravitational force provides/is the centripetal force same gravitational force (by Newton III)	B1 B1 [2]
(a)(ii)	$\omega = 2\pi / T$ $= 2\pi / (4.0 \times 365 \times 24 \times 3600)$ $= 5.0 (4.98) \times 10^{-8} \text{ rad s}^{-1}$	C1 A1 [2]
(b)(i)	(centripetal force =) $M_A d \omega^2 = M_B (2.8 \times 10^8 - d) \omega^2$ or $M_A d_A = M_B d_B$ $M_A / M_B = 3.0 = (2.8 \times 10^8 - d) / d$ $d = 7.0 \times 10^7 \text{ km}$	C1 C1 A1 [3]
(b)(ii)	$G M_A M_B / (2.8 \times 10^{11})^2 = M_A d \omega^2$ $M_B = (2.8 \times 10^{11})^2 \times d \omega^2 / G$ $= (2.8 \times 10^{11})^2 \times (7.0 \times 10^{10}) \times (4.98 \times 10^{-8})^2 / (6.67 \times 10^{-11})$ $= 2.0 \times 10^{29} \text{ kg}$	B1 C1 A1 [3]

[Total: 10]

- 1 A binary star consists of two stars A and B that orbit one another, as illustrated in Fig. 1.1.




Fig. 1.1

The stars are in circular orbits with the centres of both orbits at point P, a distance  $d$  from the centre of star A.

- (a) (i) Explain why the centripetal force acting on both stars has the same magnitude.

The <sup>Torque</sup> forces of on both stars are a couple. The <sup>of torque</sup> force is equal to the centripetal force which is the torque. [2]

- (ii) The period of the orbit of the stars about point P is 4.0 years. <sup>same.</sup> them, which are

Calculate the angular speed  $\omega$  of the stars.

$$\begin{aligned}\omega &= \frac{2\pi}{T} \\ &= \frac{2\pi}{126144000} \\ &= 4.98 \times 10^{-8} \\ &= 5.0 \times 10^{-8}\end{aligned}$$

$$\begin{aligned}T &= 4 \times 365 \times 24 \times 60 \times 60 \\ &= 126144000 \text{ s}\end{aligned}$$

$$\omega = 5.0 \times 10^{-8} \text{ rad s}^{-1} \text{ [2]}$$

Your  
Mark

1(a)(i)

1(a)(ii)

1(b)(i)

1(b)(ii)

Q1	Mark scheme	
(a)(i)	gravitational force provides/is the centripetal force same gravitational force (by Newton III)	B1 B1 [2]
(a)(ii)	$\omega = 2\pi / T$ $= 2\pi / (4.0 \times 365 \times 24 \times 3600)$ $= 5.0 \text{ (4.98)} \times 10^{-8} \text{ rad s}^{-1}$	C1 A1 [2]
(b)(i)	(centripetal force =) $M_A d \omega^2 = M_B (2.8 \times 10^8 - d) \omega^2$ or $M_A d_A = M_B d_B$ $M_A / M_B = 3.0 = (2.8 \times 10^8 - d) / d$ $d = 7.0 \times 10^7 \text{ km}$	C1 C1 A1 [3]
(b)(ii)	$GM_A M_B / (2.8 \times 10^{11})^2 = M_A d \omega^2$ $M_B = (2.8 \times 10^{11})^2 \times d \omega^2 / G$ $= (2.8 \times 10^{11})^2 \times (7.0 \times 10^{10}) \times (4.98 \times 10^{-8})^2 / (6.67 \times 10^{-11})$ $= 2.0 \times 10^{29} \text{ kg}$	B1 C1 A1 [3]
		[Total: 10]

- (b) The separation of the centres of the stars is  $2.8 \times 10^8$  km.  
The mass of star A is  $M_A$ . The mass of star B is  $M_B$ .  
The ratio  $\frac{M_A}{M_B}$  is 3.0.

(i) Determine the distance  $d$ .

$$\begin{aligned} F_{\text{grav}} &= F_{\text{cent}} \\ \frac{GM_A M_B}{r^2} &= \frac{GM_A M_B}{r^2} \\ \frac{M_A}{r^2} &= \frac{M_B}{r^2} \\ \frac{M_A}{M_B} &= \frac{r^2}{r^2} \\ \frac{M_A}{M_B} &= \frac{r^2}{r^2} \\ d &= 1.6 \times 10^8 \text{ km} \quad [3] \end{aligned}$$

- (ii) Use your answers in (a)(ii) and (b)(i) to determine the mass  $M_B$  of star B.  
Explain your working.

$$\begin{aligned} \frac{GM_A M_B}{r^2} &= \frac{GM_A M_B}{r^2} \\ \frac{F}{m \omega^2 r} &= \frac{F}{m \omega^2 r} \\ F &= m \omega^2 r \\ m &= \frac{F}{\omega^2 r} \\ M_B &= 4 \times 10^{-7} \text{ kg} \quad [3] \end{aligned}$$

[Total: 10]

Your  
Mark

1(a)(i)

1(a)(ii)

1(b)(i)

1(b)(ii)

Q1	Mark scheme	
(a)(i)	gravitational force provides/is the centripetal force same gravitational force (by Newton III)	B1 B1 [2]
(a)(ii)	$\omega = 2\pi / T$ $= 2\pi / (4.0 \times 365 \times 24 \times 3600)$ $= 5.0 (4.98) \times 10^{-8} \text{ rad s}^{-1}$	C1 A1 [2]
(b)(i)	(centripetal force =) $M_A d \omega^2 = M_B (2.8 \times 10^8 - d) \omega^2$ or $M_A d_A = M_B d_B$ $M_A / M_B = 3.0 = (2.8 \times 10^8 - d) / d$ $d = 7.0 \times 10^7 \text{ km}$	C1 C1 A1 [3]
(b)(ii)	$GM_A M_B / (2.8 \times 10^{11})^2 = M_A d \omega^2$ $M_B = (2.8 \times 10^{11})^2 \times d \omega^2 / G$ $= (2.8 \times 10^{11})^2 \times (7.0 \times 10^{10}) \times (4.98 \times 10^{-8})^2 /$ $(6.67 \times 10^{-11})$ $= 2.0 \times 10^{29} \text{ kg}$	B1  C1 A1 [3]
		[Total: 10]



Cambridge Assessment International Education  
The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA, United Kingdom  
t: +44 1223 553554 f: +44 1223 553558  
e: [info@cambridgeinternational.org](mailto:info@cambridgeinternational.org) [www.cambridgeinternational.org](http://www.cambridgeinternational.org)

Copyright © UCLES March 2018