

# Interactive Example Candidate Responses

## Paper 2 (May/June 2016), Question 4

### Cambridge International AS & A Level

### Physics 9702

In order to help us develop the highest quality resources, we are undertaking a continuous programme of review; not only to measure the success of our resources but also to highlight areas for improvement and to identify new development needs.

We invite you to complete our survey by visiting the website below. Your comments on the quality and relevance of our resources are very important to us.

[www.surveymonkey.co.uk/r/GL6ZNJB](http://www.surveymonkey.co.uk/r/GL6ZNJB)

Would you like to become a Cambridge International consultant and help us develop support materials?

Please follow the link below to register your interest.

[www.cambridgeinternational.org/cambridge-for/teachers/teacherconsultants/](http://www.cambridgeinternational.org/cambridge-for/teachers/teacherconsultants/)

Copyright © UCLES 2018

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

UCLES retains the copyright on all its publications. Registered Centres are permitted to copy material from this booklet for their own internal use. However, we cannot give permission to Centres to photocopy any material that is acknowledged to a third party, even for internal use within a Centre.

- 4 (a) By reference to the direction of the propagation of energy, state what is meant by a *longitudinal* wave and by a *transverse* wave.

longitudinal: Waves that travel parallel to direction of propagation of energy.

transverse: Waves that travel at right angles to direction of propagation of energy.

[2]

- (b) The intensity of a sound wave passing through air is given by

$$I = K \rho v f^2 A^2$$

where  $I$  is the intensity (power per unit area),  
 $K$  is a constant without units,  
 $v$  is the speed of sound,  
 $\rho$  is the density of air,  
 $f$  is the frequency of the wave  
and  $A$  is the amplitude of the wave.

Show that both sides of the equation have the same SI base units.

$$I = \frac{P}{A} \quad P = \frac{F \times d}{t} \rightarrow \frac{\text{kg ms}^{-2} \times \text{m}}{\text{s}} = \frac{\text{kg m s}^{-2} \times \text{m}}{\text{s}} = \frac{\text{kg m}^2 \text{ s}^{-2}}{\text{s}} = \text{kg m}^2 \text{ s}^{-3}$$

$$\therefore I = \frac{\text{kg m}^2 \text{ s}^{-3}}{\text{m}^2} = \text{kg s}^{-3}$$

$$v p f^2 A^2 \rightarrow (\text{ms}^{-1}) (\text{kg m}^{-3}) \times (\text{s}^{-1})^2 (\text{m})^2$$

$$\rightarrow \text{kg m}^{-2} \text{ s}^{-1} \times \text{s}^{-2} \times \text{m}^2$$

$$\rightarrow \text{kg s}^{-3}$$

$$\therefore \text{LHS} = \text{RHS}$$

[3]

Your  
Mark

4(a)

4(b)

4(c)(i)

4(c)(ii)

4(d)

Q4	Mark scheme
(a)	longitudinal: vibrations/oscillations (of the particles/wave) are parallel to the direction or in the same direction (of the propagation of energy) B1 transverse: vibrations/oscillations (of the particles/wave) are perpendicular to the direction (of the propagation of energy) B1 [2]
(b)	LHS: intensity = power / area units: $\text{kg m s}^{-2} \times \text{m} \times \text{s}^{-1} \times \text{m}^{-2}$ B1 or $\text{kg m}^2 \text{s}^{-3} \times \text{m}^{-2}$ M1 RHS: units: $\text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2$ A13 [3] LHS and RHS both $\text{kg s}^{-3}$
(c)(i)	change/difference in the <u>observed/apparent</u> frequency when the source is moving (relative to the observer) B1 [1]
(c)(ii)	wavelength increases/frequency decreases/red shift B1 [1]
(d)	observed frequency = $v f_s / (v - v_s)$ C1 $550 = (340 \times 510) / (340 - v_s)$ C1 $v_s = 25 \text{ (24.7) m s}^{-1}$ A1 [3] [Total: 10]

- (c) (i) Describe the Doppler effect.

Where the observed frequency is different from the emitted frequency, when there is motion between the observer & source. [1]

- (ii) A distant star is moving away from a stationary observer.

State the effect of the motion on the light observed from the star.

The light becomes less bright, since frequency decreases (observed), and

..... [1]

- (d) A car travels at a constant speed towards a stationary observer. The horn of the car sounds at a frequency of 510 Hz and the observer hears a frequency of 550 Hz. The speed of sound in air is  $340 \text{ m s}^{-1}$ .

$s = 340$

Calculate the speed of the car.

$$550 = \frac{510 \times 340}{(340 - v)}$$

$$181,500 - 550v = 168,300$$

$$550v = 13,200$$

$$v = 24 \text{ m s}^{-1}$$

speed =  $24 \text{ m s}^{-1}$   $\text{m s}^{-1}$  [3]

[Total: 10]

Your  
Mark

4(a)

4(b)

4(c)(i)

4(c)(ii)

4(d)

Q4	Mark scheme
(a)	longitudinal: vibrations/oscillations (of the particles/wave) are parallel to the direction or in the same direction (of the propagation of energy) B1 transverse: vibrations/oscillations (of the particles/wave) are perpendicular to the direction (of the propagation of energy) B1 [2]
(b)	LHS: intensity = power / area units: $\text{kg m s}^{-2} \times \text{m} \times \text{s}^{-1} \times \text{m}^{-2}$ or $\text{kg m}^2 \text{s}^{-3} \times \text{m}^{-2}$ B1 RHS: units: $\text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2$ M1 LHS and RHS both $\text{kg s}^{-3}$ A13 [3]
(c)(i)	change/difference in the <u>observed/apparent</u> frequency when the source is moving (relative to the observer) B1 [1]
(c)(ii)	wavelength increases/frequency decreases/red shift B1 [1]
(d)	observed frequency = $vf_s / (v - v_s)$ C1 $550 = (340 \times 510) / (340 - v_s)$ C1 $v_s = 25 (24.7) \text{ m s}^{-1}$ A1 [3] [Total: 10]

- 4 (a) By reference to the direction of the propagation of energy, state what is meant by a *longitudinal* wave and by a *transverse* wave.

longitudinal: A wave in which the particle moves parallel to the propagation of energy is known as longitudinal wave.

transverse: A wave in which the particle of the motion moves perpendicular to the direction of motion is known as a transverse wave.

[2]

- (b) The intensity of a sound wave passing through air is given by

$$I = K \rho v f^2 A^2$$

where  $I$  is the intensity (power per unit area),

$K$  is a constant without units,

$v$  is the speed of sound,

$\rho$  is the density of air,

$f$  is the frequency of the wave

and  $A$  is the amplitude of the wave.

$$\frac{\text{mass}}{\text{vol}} = \text{kg m}^{-3}$$

Show that both sides of the equation have the same SI base units.

$$\begin{aligned} \rightarrow \frac{\text{kg} \times \text{m}^2 \times \text{s}^{-2} \times \text{m}^2}{\text{s}} &= K \times \text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2 \\ \text{kg} \times \text{s}^{-3} &= K \times \text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2 \\ \text{kg s}^{-3} &= K \times \text{m}^0 \times \text{kg} \times \text{s}^{-3} \\ \text{kg s}^{-3} &= K \text{ kg s}^{-3} \end{aligned}$$

$$H_2 = f \lambda = \frac{1}{T}$$

$$\approx \text{S s}^{-1}$$

$$E = P \times t$$

$$P = \frac{W}{t}$$

$$\begin{aligned} f \times d &= \text{ms}^{-2} \\ &= \frac{\text{kg} \times \text{a} \times \text{m}}{\text{s}} \end{aligned}$$

[3]

Your  
Mark

4(a)

4(b)

4(c)(i)

4(c)(ii)

4(d)

Q4	Mark scheme
(a)	<p>longitudinal: vibrations/oscillations (of the particles/wave) are parallel to the direction or in the same direction (of the propagation of energy) B1</p> <p>transverse: vibrations/oscillations (of the particles/wave) are perpendicular to the direction (of the propagation of energy) B1 [2]</p>
(b)	<p>LHS: intensity = power / area units: <math>\text{kg m s}^{-2} \times \text{m} \times \text{s}^{-1} \times \text{m}^{-2}</math> B1 or <math>\text{kg m}^2 \text{s}^{-3} \times \text{m}^{-2}</math> M1 RHS: units: <math>\text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2</math> A13 [3] LHS and RHS both <math>\text{kg s}^{-3}</math></p>
(c)(i)	<p>change/difference in the <u>observed/apparent</u> frequency when the source is moving (relative to the observer) B1 [1]</p>
(c)(ii)	<p>wavelength increases/frequency decreases/red shift B1 [1]</p>
(d)	<p>observed frequency = <math>v f_s / (v - v_s)</math> C1 <math>550 = (340 \times 510) / (340 - v_s)</math> C1 <math>v_s = 25 \text{ (24.7) m s}^{-1}</math> A1 [3] [Total: 10]</p>

- (c) (i) Describe the Doppler effect.

*It The observed frequency is always different to the frequency emitted when source and sound are in a relative motion.* [1]

- (ii) A distant star is moving away from a stationary observer.

State the effect of the motion on the light observed from the star,

*The wavelength and the frequency has now been changed so the motion will also change.* [1]

- (d) A car travels at a constant speed towards a stationary observer. The horn of the car sounds at a frequency of 510 Hz and the observer hears a frequency of 550 Hz. The speed of sound in air is  $340 \text{ m s}^{-1}$ .

Calculate the speed of the car.

$$f = 510$$

$$510 = 550 \times \left( \frac{340}{340 - v_{\text{car}}} \right)$$

$$0.927 (340 - v_{\text{car}}) = 340$$

$$340 - v_{\text{car}} = 366.67$$

$$v_{\text{car}} = 26.67$$

speed = ~~266~~ 26.67  $\text{m s}^{-1}$  [3]

[Total: 10]

Your  
Mark

4(a)

4(b)

4(c)(i)

4(c)(ii)

4(d)

Q4	Mark scheme
(a)	longitudinal: vibrations/oscillations (of the particles/wave) are parallel to the direction or in the same direction (of the propagation of energy) B1 transverse: vibrations/oscillations (of the particles/wave) are perpendicular to the direction (of the propagation of energy) B1 [2]
(b)	LHS: intensity = power / area units: $\text{kg m s}^{-2} \times \text{m} \times \text{s}^{-1} \times \text{m}^{-2}$ or $\text{kg m}^2 \text{s}^{-3} \times \text{m}^{-2}$ B1 RHS: units: $\text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2$ M1 LHS and RHS both $\text{kg s}^{-3}$ A13 [3]
(c)(i)	change/difference in the <u>observed/apparent</u> frequency when the source is moving (relative to the observer) B1 [1]
(c)(ii)	wavelength increases/frequency decreases/red shift B1 [1]
(d)	observed frequency = $vf_s / (v - v_s)$ C1 $550 = (340 \times 510) / (340 - v_s)$ C1 $v_s = 25 (24.7) \text{ m s}^{-1}$ A1 [3] [Total: 10]

- 4 (a) By reference to the direction of the propagation of energy, state what is meant by a *longitudinal* wave and by a *transverse* wave.

longitudinal: The energy is ~~not~~ parallel to the direction of the propagation, such as a sound wave.

transverse: The energy is perpendicular to the direction of the propagation, such as in a guitar string.

[2]

- (b) The intensity of a sound wave passing through air is given by

$$I = K \rho v f^2 A^2$$

where  $I$  is the intensity (power per unit area),

$K$  is a constant without units,

$v$  is the speed of sound,

$\rho$  is the density of air,

$f$  is the frequency of the wave

and  $A$  is the amplitude of the wave.

so that both sides of the equation have the same SI base units.

$$I = K \rho v f^2 A^2$$

$$\frac{\text{m}^2 \text{s}^{-4}}{\text{m}^2} = \text{m}^3 \text{s}^{-1} \text{kg m}^{-3} \text{s}^{-3} \text{m}^2$$

$$\text{kg s}^{-4} = \text{kg s}^{-4}$$

Thus proved

$$\text{power} = \frac{f \times d}{t}$$

$$\begin{aligned} \text{power} &= \frac{\text{m ad}}{t} \\ &= \frac{\text{kg m s}^{-2} \text{m}}{s} \\ &= \frac{\text{kg m}^2 \text{s}^{-2}}{s} \\ &= \text{kg m}^2 \text{s}^{-3} \end{aligned}$$

$$-v = \frac{\text{m}}{\text{s}} = \text{m s}^{-1}$$

$$-\rho = \frac{\text{m}}{\text{m}^3} = \frac{\text{kg}}{\text{m}^3} = \text{kg m}^{-3}$$

$$-f = \frac{1}{\text{s}} = \text{s}^{-1}$$

$$-A = \text{m}^2$$

[3]

Your  
Mark

4(a)

4(b)

4(c)(i)

4(c)(ii)

4(d)

Q4	Mark scheme
(a)	<p>longitudinal: vibrations/oscillations (of the particles/wave) are parallel to the direction or in the same direction (of the propagation of energy) B1</p> <p>transverse: vibrations/oscillations (of the particles/wave) are perpendicular to the direction (of the propagation of energy) B1 [2]</p>
(b)	<p>LHS: intensity = power / area units: <math>\text{kg m s}^{-2} \times \text{m} \times \text{s}^{-1} \times \text{m}^{-2}</math> B1 or <math>\text{kg m}^2 \text{s}^{-3} \times \text{m}^{-2}</math> M1 RHS: units: <math>\text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2</math> A13 [3] LHS and RHS both <math>\text{kg s}^{-3}</math></p>
(c)(i)	change/difference in the <u>observed/apparent</u> frequency when the source is moving (relative to the observer) B1 [1]
(c)(ii)	wavelength increases/frequency decreases/red shift B1 [1]
(d)	<p>observed frequency = <math>v f_s / (v - v_s)</math> C1  <math>550 = (340 \times 510) / (340 - v_s)</math> C1  <math>v_s = 25 \text{ (24.7) m s}^{-1}</math> A1 [3]</p>
[Total: 10]	



(c) (i) Describe the Doppler effect.

The change in apparent frequency due to the change in movement of the source or observer. [1]

(ii) A distant star is moving away from a stationary observer.

State the effect of the motion on the light observed from the star.

The apparent frequency would decrease. [1]

(d) A car travels at a constant speed towards a stationary observer. The horn of the car sounds at a frequency of 510 Hz and the observer hears a frequency of 550 Hz. The speed of sound in air is  $340 \text{ m s}^{-1}$ .

Calculate the speed of the car.

$$f_o = \left( \frac{v_w}{v_w - v_s} \right) f_s$$

$$550 = \left( \frac{340}{340 - v_s} \right) 510$$

$$40 = \frac{340}{340 - v_s}$$

$$340 = 13600 - 40v_s$$

$$40v_s = 13600 - 3400$$

$$v_s = 331.5$$

speed = 331.5  $\text{m s}^{-1}$  [3]

[Total: 10]

Your  
Mark

4(a)

4(b)

4(c)(i)

4(c)(ii)

4(d)

Q4	Mark scheme
(a)	<p>longitudinal: vibrations/oscillations (of the particles/wave) are parallel to the direction or in the same direction (of the propagation of energy) B1</p> <p>transverse: vibrations/oscillations (of the particles/wave) are perpendicular to the direction (of the propagation of energy) B1 [2]</p>
(b)	<p>LHS: intensity = power / area units: <math>\text{kg m s}^{-2} \times \text{m} \times \text{s}^{-1} \times \text{m}^{-2}</math> or <math>\text{kg m}^2 \text{s}^{-3} \times \text{m}^{-2}</math> B1</p> <p>RHS: units: <math>\text{m s}^{-1} \times \text{kg m}^{-3} \times \text{s}^{-2} \times \text{m}^2</math> M1</p> <p>LHS and RHS both <math>\text{kg s}^{-3}</math> A13 [3]</p>
(c)(i)	<p>change/difference in the <u>observed/apparent</u> frequency when the source is moving (relative to the observer) B1 [1]</p>
(c)(ii)	<p>wavelength increases/frequency decreases/red shift B1 [1]</p>
(d)	<p>observed frequency = <math>v f_s / (v - v_s)</math> C1</p> <p><math>550 = (340 \times 510) / (340 - v_s)</math> C1</p> <p><math>v_s = 25 \text{ (24.7) m s}^{-1}</math> A1 [3]</p> <p>[Total: 10]</p>



Cambridge Assessment International Education  
The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA, United Kingdom  
t: +44 1223 553554 f: +44 1223 553558  
e: [info@cambridgeinternational.org](mailto:info@cambridgeinternational.org) [www.cambridgeinternational.org](http://www.cambridgeinternational.org)

Copyright © UCLES March 2018