



2.1 Particles in the atom

KC




2.2 The nucleus of the atom

KC




1.1 Relative masses of atoms and molecules

KC




2.3 Electrons: energy levels, atomic orbitals, ionisation energy, electron affinity

KC


 6.022×10^{23} 1 MOLE


1.2 The mole and the avogadro constant

KC


1.3 The determination of relative atomic masses, A_r

KC



1.4 The calculation of empirical and molecular formula

KC



1.5 Reacting masses and volumes (of solutions and gases)


KC

Formula mass, relative atomic mass, and moles


Subatomic particles


2


Bonding and structure



Controlling reactions 1










6

Acids and bases









STRONG ACID



WEAK ACID




$$\text{pH} = -\log[\text{H}^+(\text{aq})]$$



7.2

Ionic equilibria






$$F = Le \quad -1.602 \times 10^{-19} \text{ C}$$


+ anode metal OR H_2
 - cathode halogen OR O_2

6.2

Electrolysis




6.3

Standard electrode potential E° ; standard cell potentials E°_{cell} and the Nernst equation


12.2

General characteristic chemical properties of the first set of transition elements, titanium to copper




6.4

Batteries and fuel cells



12.1

General physical properties of the first set of transition elements, titanium to copper




12.3

Colour of complexes



ligand substitution reaction




$$K_{\text{stab}} = \frac{[\text{products}]}{[\text{reactants}]} = \frac{[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2^{2+}]}{[\text{Cu}(\text{H}_2\text{O})_6^{2+}] [\text{NH}_3]^4}$$


12.5


Stability constants, K_{stab} 

8


Controlling reactions 2








22.3 Mass spectrometry KC



22.1 Chromatography KC

Structure determination


Separating mixtures




$$K_{pc} = \frac{[X \text{ in solvent 1}]}{[X \text{ in solvent 2}]}$$

- X (solute)
- solvent 1
- solvent 2


7.3 Partition coefficients KC



22.4 Carbon-13 NMR spectroscopy KC



22.5 Proton (1H) NMR spectroscopy KC



NMR