

Interactive Example Candidate Responses

Paper 5 (May / June 2016), Question 2

Cambridge IGCSE™
Physics 0625



In order to help us develop the highest quality resources, we are undertaking a continuous programme of review; not only to measure the success of our resources but also to highlight areas for improvement and to identify new development needs.

We invite you to complete our survey by visiting the website below. Your comments on the quality and relevance of our resources are very important to us.

www.surveymonkey.co.uk/r/GL6ZNJB

Would you like to become a Cambridge International consultant and help us develop support materials?

Please follow the link below to register your interest.

www.cambridgeinternational.org/cambridge-for/teachers/teacherconsultants/

Copyright © UCLES 2017

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

UCLES retains the copyright on all its publications. Registered Centres are permitted to copy material from this booklet for their own internal use. However, we cannot give permission to Centres to photocopy any material that is acknowledged to a third party, even for internal use within a Centre.

2 In this experiment, you will investigate the cooling of water.

- (a) • Pour 100 cm³ of the hot water provided into beaker A.

- Measure the temperature θ_H of the water in beaker A.

$$\theta_H = \dots\dots\dots 86^\circ\text{C}$$

- Pour 100 cm³ of the cold water provided into beaker B.

- Measure the temperature θ_C of the water in beaker B.

$$\theta_C = \dots\dots\dots 30^\circ\text{C}$$

- Calculate the average temperature θ_{AV} using the equation $\theta_{AV} = \frac{\theta_H + \theta_C}{2}$.

$$\begin{array}{r} 86 + 30 \\ \hline 2 \\ \hline \theta_{AV} = \dots\dots\dots 58^\circ\text{C} \end{array} \quad [3]$$

- (b) Add the water from beaker B to the hot water in beaker A. Stir briefly.

Measure the temperature θ_M of the mixture.

$$\theta_M = \dots\dots\dots 51^\circ\text{C} \quad [1]$$

- (c) State one precaution that you took to ensure that the temperature readings are as reliable as possible.

Make sure that I take the readings from eye level to prevent parallax error. [1]

Select
page

Your
Mark

2(a)

2(b)

2(c)

2(d)(i)

2(d)(ii)

2(d)(iii)

2(d)(iv)


Q2	Mark scheme
(a)	θ_H 60 – 100 θ_C 10 – 40 and θ_{AV} correct Unit °C
(b)	θ_M between θ_H and θ_C
(c)	Perpendicular viewing of scale OR wait until temperature stops rising OR carry out without undue delay between parts
(d)(i)	Correct diagram with lid Insulation placed round beaker
(d)(ii)	Sensible series of values with θ_M between θ_H and θ_C
(d)(iii)	Statement and justification to match results
(d)(iv)	Two from: Room temperature (or other environmental condition) Temperature of cold water Temperature of hot water Volumes of water Size/shape/material/surface area of beaker

(d) Empty both beakers.

You are provided with

- a lid, with a hole for the thermometer,
- some insulating material,
- two elastic bands.

(i) In the space below, draw a labelled diagram to show how you will use these items to reduce the loss of thermal energy when the procedure is repeated.



[2]

(ii) Using the improvements shown in your diagram, repeat the procedure in parts (a) and (b).

$\theta_H = \dots\dots\dots 62^\circ \quad 73^\circ C$
 $\theta_C = \dots\dots\dots 31^\circ C$
 $\theta_{AV} = \dots\dots\dots 52^\circ C$
 $\theta_M = \dots\dots\dots 50^\circ C$

[1]

(iii) Comment on whether the improvements made to the apparatus have significantly changed the value of the temperature θ_M . Use your results to justify your answer.

~~No~~ The value of θ_M has not significantly changed because there is only a $1^\circ C$ difference between both experiments.

[1]

(iv) Suggest two conditions that should be kept constant for all parts of this experiment.

1. The amount of water used
2. The ^{room} external ^{temperature} environment must be maintained

[2]

[Total: 11]

Your
Mark

2(a)

2(b)

2(c)

2(d)(i)

2(d)(ii)

2(d)(iii)

2(d)(iv)

Q2 Mark scheme

(a)	θ_H 60 – 100 θ_C 10 – 40 and θ_{AV} correct Unit $^\circ C$
(b)	θ_M between θ_H and θ_C
(c)	Perpendicular viewing of scale OR wait until temperature stops rising OR carry out without undue delay between parts
(d)(i)	Correct diagram with lid Insulation placed round beaker
(d)(ii)	Sensible series of values with θ_M between θ_H and θ_C
(d)(iii)	Statement and justification to match results
(d)(iv)	Two from: Room temperature (or other environmental condition) Temperature of cold water Temperature of hot water Volumes of water Size/shape/material/surface area of beaker

2 In this experiment, you will investigate the cooling of water.

- (a) • Pour 100 cm³ of the hot water provided into beaker A.

- Measure the temperature θ_H of the water in beaker A.

$$\theta_H = 66^\circ\text{C}$$

- Pour 100 cm³ of the cold water provided into beaker B.

- Measure the temperature θ_C of the water in beaker B.

$$\theta_C = 33^\circ\text{C}$$

- Calculate the average temperature θ_{AV} using the equation $\theta_{AV} = \frac{\theta_H + \theta_C}{2}$.

$$\frac{66 + 33}{2} = 49.5$$

$$\theta_{AV} = 49^\circ\text{C}$$

- (b) Add the water from beaker B to the hot water in beaker A. Stir briefly.

Measure the temperature θ_M of the mixture.

$$\theta_M = 47^\circ\text{C}$$

- (c) State one precaution that you took to ensure that the temperature readings are as reliable as possible.

I tried to avoid parallax error when pouring water into the measuring cylinder.

Select page

Your Mark

2(a)

2(b)

2(c)

2(d)(i)

2(d)(ii)

2(d)(iii)

2(d)(iv)


Q2	Mark scheme
(a)	θ_H 60 – 100 θ_C 10 – 40 and θ_{AV} correct Unit °C
(b)	θ_M between θ_H and θ_C
(c)	Perpendicular viewing of scale OR wait until temperature stops rising OR carry out without undue delay between parts
(d)(i)	Correct diagram with lid Insulation placed round beaker
(d)(ii)	Sensible series of values with θ_M between θ_H and θ_C
(d)(iii)	Statement and justification to match results
(d)(iv)	Two from: Room temperature (or other environmental condition) Temperature of cold water Temperature of hot water Volumes of water Size/shape/material/surface area of beaker

(d) Empty both beakers.

You are provided with

- a lid, with a hole for the thermometer,
- some insulating material,
- two elastic bands.

(i) In the space below, draw a labelled diagram to show how you will use these items to reduce the loss of thermal energy when the procedure is repeated.



[2]

(ii) Using the improvements shown in your diagram, repeat the procedure in parts (a) and (b).

$$\frac{69 + 33}{2} = 51$$

$$\theta_H = 69^\circ\text{C}$$

$$\theta_C = 33^\circ\text{C}$$

$$\theta_{AV} = 51^\circ\text{C}$$

$$\theta_M = 50^\circ\text{C}$$

[1]

(iii) Comment on whether the improvements made to the apparatus have significantly changed the value of the temperature θ_M . Use your results to justify your answer.

Yes they have. θ_M without insulation was 47°C and θ_M with insulation was 50°C . Heat loss with insulation is more.

(iv) Suggest two conditions that should be kept constant for all parts of this experiment.

1. The initial ~~room~~ room temperature
2. The volume of water added

[2]

[Total: 11]

Your
Mark

2(a)

2(b)

2(c)

2(d)(i)

2(d)(ii)

2(d)(iii)

2(d)(iv)

Q2 Mark scheme

(a)	θ_H 60 – 100 θ_C 10 – 40 and θ_{AV} correct Unit $^\circ\text{C}$
(b)	θ_M between θ_H and θ_C
(c)	Perpendicular viewing of scale OR wait until temperature stops rising OR carry out without undue delay between parts
(d)(i)	Correct diagram with lid Insulation placed round beaker
(d)(ii)	Sensible series of values with θ_M between θ_H and θ_C
(d)(iii)	Statement and justification to match results
(d)(iv)	Two from: Room temperature (or other environmental condition) Temperature of cold water Temperature of hot water Volumes of water Size/shape/material/surface area of beaker

2 In this experiment, you will investigate the cooling of water.

- (a) • Pour 100 cm³ of the hot water provided into beaker A.

- Measure the temperature θ_H of the water in beaker A.

$$\theta_H = 78^\circ$$

- Pour 100 cm³ of the cold water provided into beaker B.

- Measure the temperature θ_C of the water in beaker B.

$$\theta_C = 32^\circ$$

- Calculate the average temperature θ_{AV} using the equation $\theta_{AV} = \frac{\theta_H + \theta_C}{2}$.

$$\theta_{AV} = \frac{78 + 32}{2}$$

$$\theta_{AV} = 55$$

$$\theta_{AV} = 55^\circ$$

[3]

- (b) Add the water from beaker B to the hot water in beaker A. Stir briefly.

Measure the temperature θ_M of the mixture.

$$\theta_M = 52^\circ$$

[1]

- (c) State one precaution that you took to ensure that the temperature readings are as reliable as possible.

keep room temperature constant. Use the same volume of water for both hot and cold water

[1]

Select
page

Your
Mark

2(a)

2(b)

2(c)

2(d)(i)

2(d)(ii)

2(d)(iii)

2(d)(iv)


Q2	Mark scheme
(a)	θ_H 60 – 100 θ_C 10 – 40 and θ_{AV} correct Unit °C
(b)	θ_M between θ_H and θ_C
(c)	Perpendicular viewing of scale OR wait until temperature stops rising OR carry out without undue delay between parts
(d)(i)	Correct diagram with lid Insulation placed round beaker
(d)(ii)	Sensible series of values with θ_M between θ_H and θ_C
(d)(iii)	Statement and justification to match results
(d)(iv)	Two from: Room temperature (or other environmental condition) Temperature of cold water Temperature of hot water Volumes of water Size/shape/material/surface area of beaker

(d) Empty both beakers.

You are provided with

- a lid, with a hole for the thermometer,
- some insulating material,
- two elastic bands.

(i) In the space below, draw a labelled diagram to show how you will use these items to reduce the loss of thermal energy when the procedure is repeated.



[2]

(ii) Using the improvements shown in your diagram, repeat the procedure in parts (a) and (b).

$\theta_H = 77^\circ$
 $\theta_C = 32^\circ$
 $\theta_{AV} = 54.5^\circ$
 $\theta_M = 56^\circ$

[1]

(iii) Comment on whether the improvements made to the apparatus have significantly changed the value of the temperature θ_M . Use your results to justify your answer.

$\frac{56 - 52}{56} \times 100 = 7.14\%$. Yes it has. No it has.
 56 Yes it has changed the value, because it
 has changed increased by 4° from 52° to 56° .

[1]

(iv) Suggest two conditions that should be kept constant for all parts of this experiment.

1. Initial temperature
2. room temperature

[2]

[Total: 11]

Your
Mark

2(a)

2(b)

2(c)

2(d)(i)

2(d)(ii)

2(d)(iii)

2(d)(iv)

Q2	Mark scheme
(a)	θ_H 60 – 100 θ_C 10 – 40 and θ_{AV} correct Unit $^\circ\text{C}$
(b)	θ_M between θ_H and θ_C
(c)	Perpendicular viewing of scale OR wait until temperature stops rising OR carry out without undue delay between parts
(d)(i)	Correct diagram with lid Insulation placed round beaker
(d)(ii)	Sensible series of values with θ_M between θ_H and θ_C
(d)(iii)	Statement and justification to match results
(d)(iv)	Two from: Room temperature (or other environmental condition) Temperature of cold water Temperature of hot water Volumes of water Size/shape/material/surface area of beaker

Cambridge Assessment International Education
The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA, United Kingdom
t: +44 1223 553554
e: info@cambridgeinternational.org www.cambridgeinternational.org

Copyright © UCLES September 2017